

nant magnetic carrier in the sheeted dike basalts, did not form simply by oxidation-exsolution, as has commonly been assumed (6, 12, 26). The natural remanent magnetization (NRM) of the sheeted-dike basalts was presumably acquired by the single-domain Ti-bearing magnetite (the host after exsolution) during initial cooling. This was a thermoremanent magnetization (TRM) and was then modified by chemical remanent magnetization of recrystallized end-member magnetite during hydrothermal alteration near the spreading center. The initial cooling and hydrothermal alteration appear to have taken place soon after the intrusion of basalts; therefore, the NRM reflects the original geomagnetic field direction (6).

The thickness of the source layers responsible for the sea-floor magnetic anomalies has long been debated and has been estimated as extending from the uppermost 500 to 1000 m (pillow basalts, layer 2A) of the oceanic crust to depths of ~8 km (essentially the entire oceanic crust) (3, 7, 27, 28). The results of studies of magnetic properties of sheeted dike basalts recovered from DSDP drill holes suggest that the sheeted dike complex (layer 2B) contributes significantly to sea-floor magnetic anomalies (6, 26, 29). However, magnetic data from ocean gabbros indicate that the linear magnetic anomalies originated partly in the gabbro layer (layer 3) (29-31). We have shown that single-domain, end-member magnetite, an efficient and stable carrier of TRM, is responsible for the magnetic properties in the upper levels (depths of ~30 m to at least ~1125 m within the igneous basement) of the sheeted dike complex at site 504B. The resultant NRM intensity of the sheeted dike basalts is on the same order as that of the pillow basalts at site 504B (6, 8, 12). We therefore conclude that the upper sheeted dike basalts from DSDP hole 504B are a significant source of sea-floor magnetic anomalies.

REFERENCES AND NOTES

1. F. J. Vine and D. H. Matthews, *Nature* 190, 947 (1963).
2. F. J. Vine and J. T. Wilson, *Science* 150, 485 (1965).
3. C. G. A. Harrison, in *The Oceanic Lithosphere*, vol. 7 of *The Sea*, C. Emiliani, Ed. (Wiley, New York, 1981), pp. 219-239.
4. H. P. Johnson and J. M. Hall, *Geophys. J. R. Astron. Soc.* 52, 45 (1976).
5. N. Peterson, P. Elanach, U. Bleil, in *Deep Drilling Results in the Atlantic Ocean: Ocean Crust*, M. Tahani, C. G. Harrison, D. E. Hayes, Eds. (American Geophysical Union, Washington, DC, 1979), vol. 2, pp. 169-202.
6. G. M. Smith and S. K. Banerjee, *J. Geophys. Res.* 91, 10337 (1986).
7. H. P. Johnson and T. Alwater, *Geol. Soc. Am. Bull.* 88, 637 (1977).
8. T. Furuta, *Int. Rep. Deep Sea Drill. Prof.* 69, 711 (1983).
9. J. B. O'Donovan and W. O'Reilly, *ibid.*, p. 721.
10. D. M. Peckersky, V. Tikhonov, N. N. Pertsev, *ibid.*, p. 705.
11. G. M. Smith and S. K. Banerjee, *ibid.* 83, 347 (1985).
12. D. Facey, J. Housden, W. O'Reilly, *ibid.*, p. 339.
13. H. Kinoshita, T. Furuta, H. Kawahata, *ibid.*, p. 331.
14. Oxidation-exsolution is defined as oxidation reactions with oxygen partitioning into titanomagnetite but giving rise to well-oriented lamellar textures that are commonly caused by exsolution.
15. A Philips CM-12 scanning transmission electron microscope equipped with a Kevex (Chesham, United Kingdom) quantum detector was used. The methods for specimen preparation and STEM quantitative chemical analyses are described in (16).
16. Y.-H. Shau, H.-Y. Yang, D. R. Peacor, *Am. Mineral.* 76, 1208 (1991).
17. J. C. Alt, J. Honnorez, C. Laverne, R. Emmertmann, *J. Geophys. Res.* 91, 10309 (1986).
18. Y.-H. Shau and D. R. Peacor, *Contrib. Mineral. Petrol.* 112, 119 (1992).
19. S. E. Haggerty, in *Oxide Minerals, Reviews in Mineralogy*, D. Rumble, Ed. (Mineralogical Society of America, Washington, DC, 1978), vol. 3, pp. Hg101-Hg300.
20. The minor amounts of Ti, Si, and Ca cannot account for intimately intergrown sphene and magnetite lamellae in the relatively thick areas. Thus, the lamellar voids also exist in the thick areas; they are real texture and not an artifact caused by ion milling.
21. D. H. Lindsay, *Am. Mineral.* 68, 769 (1981).
22. G. D. Price, *ibid.*, p. 751.
23. P. P. K. Smith, *ibid.* 68, 1038 (1980).
24. M. E. Evans and M. L. Wayman, *Can. J. Earth Sci.* 9, 671 (1972).
25. R. F. Butler and S. K. Banerjee, *J. Geophys. Res.* 80, 4049 (1975).
26. J. E. Parise and H. P. Johnson, *ibid.* 96, 11703 (1991).
27. S. K. Banerjee, *Tectonophysics* 108, 15 (1984).
28. C. G. A. Harrison, *Annu. Rev. Earth Planet. Sci.* 15, 505 (1987).
29. D. J. Dunlop and M. Prévot, *Geophys. J. R. Astron. Soc.* 68, 763 (1982).
30. E. Kikawa and K. Ozawa, *Science* 258, 706 (1992).
31. D. V. Kent, B. M. Honnorez, N. D. Opdyke, P. J. Fox, *Geophys. J. R. Astron. Soc.* 53, 513 (1978).
32. Analyses were calculated as end-member components (all in mole percent): magnetite, 100% Fe₃O₄; ulvöspinel, 88.8% Fe₂TiO₄, 9.2% Fe₃O₄, and 4.0% Mn₂TiO₄; ilmenite, 80.6% FeTiO₃, and 9.4% MnTiO₃. All components are calculated after subtraction of the Ca, Ti, and Si that were present as a result of contamination by sphene.
33. We thank H. E. Robertson for providing the basalt samples, R. Van der Voo and two anonymous reviewers for comments, and J. C. Alt and D. Suk for many helpful discussions.

26 January 1993; accepted 11 May 1993

Induction of Apoptosis by the Low-Affinity NGF Receptor

Shahrooz Rabizadeh, Justin Oh, Li-tao Zhong, Jie Yang, Catherine M. Bitler, Larry L. Butcher, Dale E. Bredesen*

Nerve growth factor (NGF) binding to cellular receptors is required for the survival of some neural cells. In contrast to Trk A, the high-affinity NGF receptor that transduces NGF signals for survival and differentiation, the function of the low-affinity NGF receptor, p75^{NGFR}, remains uncertain. Expression of p75^{NGFR} induced neural cell death constitutively when p75^{NGFR} was unbound; binding by NGF or monoclonal antibody, however, inhibited cell death induced by p75^{NGFR}. Thus, expression of p75^{NGFR} may explain the dependence of some neural cells on NGF for survival. These findings also suggest that p75^{NGFR} has some functional similarities to other members of a superfamily of receptors that include tumor necrosis factor receptors, Fas (Apo-1), and CD40.

Growth factors such as NGF enhance the survival of cells displaying the appropriate receptors. The effects of NGF are mediated at least in part by Trk A, the high-affinity NGF receptor, which is a tyrosine kinase (1). The low-affinity NGF receptor, p75^{NGFR}, is a receptor of incompletely characterized function: p75^{NGFR} has been shown to increase the affinity of Trk A for NGF (1) and to enhance the specificity of

the Trk family of receptors for neurotrophins (2). p75^{NGFR} has some sequence similarity to the tumor necrosis factor receptors [TNFR I (3) and TNFR II (4)], the human cell surface antigen Fas (Apo-1) (5), and the B cell antigen CD40 (6), all of which mediate cell death. In the case of TNFR I and Fas, binding of the receptor by ligand or antibody initiates cell death. In the case of CD40, however, binding by monoclonal antibody (mAb) or ligand inhibits cell death (6). Thus, some cells expressing CD40 are dependent on ligand or mAb binding for survival. Because of structural and functional analogies between the CD40 and p75^{NGFR} systems, the possibility that p75^{NGFR} serves as a constitutive cell death-promoting molecule that is inhibited by NGF binding was evaluated.

We expressed p75^{NGFR} in temperature-

S. Rabizadeh, L. Zhong, J. Yang, Department of Neurology, University of California, Los Angeles, CA 90024.


J. Oh and L. L. Butcher, Department of Psychology, University of California, Los Angeles, CA 90024.

C. M. Bitler, Stanford Research Institute, Menlo Park, CA 94025.

D. E. Bredesen, Department of Neurology and Molecular Biology Institute, University of California, Los Angeles, CA 90024.

*To whom correspondence should be addressed.

Fig. 1. Expression of rat $p75^{NGFR}$ by transfected conditionally immortalized neural cells (28). (A) Northern blot demonstrating lack of Trk A and $p75^{NGFR}$ expression by CSM 14.1 cells. $p75^{NGFR}$ expression was demonstrated after transfection by pBabe-puro- $p75^{NGFR}$. Lane 1, CSM 14.1 transfected with pBabe-puro- $p75^{NGFR}$ and grown in serum-containing medium without NGF; lane 2, CSM 14.1 transfected with pBabe-puro- $p75^{NGFR}$ and grown in medium with serum and NGF (2 nM); lane 3, CSM 14.1 transfected with pBabe-puro and grown in serum-containing medium without NGF; lane 4, CSM 14.1 transfected with pBabe-puro and grown in medium with serum and NGF (2 nM); and lane 5, PC12 control. Note that the endogenous transcript in PC12 cells [3.7 kb (29)] is shorter than the transcript in the pBabe-puro- $p75^{NGFR}$ -



transfected cells (predicted to be 4.3 kb), and that treatment of the CSM 14.1 cells with NGF did not result in $p75^{NGFR}$ expression (lanes 2 and 4). Lanes 1 through 4 contained 25 μ g of total RNA; lane 5 contained 10 μ g of total RNA. (B) Immunocytochemistry of CSM 14.1 cells transfected with pBabe-puro- $p75^{NGFR}$ ($85 \pm 11\%$ of cells expressed $p75^{NGFR}$). (C) Immunocytochemistry of CSM 14.1 cells transfected with pBabe-puro ($0.6 \pm 0.5\%$ of the cells expressed $p75^{NGFR}$). Magnification, $\times 400$.

sensitive immortalized neural cells (7) by means of a retroviral vector, pBabe-puro- $p75^{NGFR}$ (8) (Fig. 1). Control cells transfected with pBabe-puro expressed neither $p75^{NGFR}$ nor Trk A (Fig. 1). In cells cultured in medium containing serum, expression of $p75^{NGFR}$ had no effect on cell death, but when serum was withdrawn to induce apoptosis (9), expression of $p75^{NGFR}$ led to an increase in neural cell death (Fig. 2). However, if NGF (5 nM) was added, not only was the negative effect on cell survival suppressed, but the cells had a death rate less than that of control cells transfected with the identical vector lacking the $p75^{NGFR}$ sequence (Fig. 2). Binding of $p75^{NGFR}$ by a mAb also suppressed the enhancement of neural cell death by $p75^{NGFR}$, but led to less improvement of cell survival than did NGF (Fig. 2). Addition of a control mAb did not affect cell survival (Fig. 2). Neither NGF nor mAb affected survival of the control cells (Fig. 2).

We demonstrated that the type of cell death induced by $p75^{NGFR}$ was apoptotic by expressing $p75^{NGFR}$ in the R2 cell line, a conditionally immortalized cerebellar neural line (10) that, in the absence of $p75^{NGFR}$ expression, does not undergo apoptosis in serum-free medium. As shown in Fig. 3, expression of $p75^{NGFR}$ by the R2 cell line led to virtually complete cell death in serum-free medium, with the nuclear fragmentation, chromatin condensation, and homogeneous nuclear staining that are characteristic of apoptosis but not necrosis (11). Control R2 transfectants survived well in serum-free medium (Fig. 3).

It was possible that the mediation of neural cell death by $p75^{NGFR}$ might have been a result of the vector-driven expression of $p75^{NGFR}$ in neural cells that do not express endogenous $p75^{NGFR}$. Therefore, PC12 pheochromocytoma cells, which express $p75^{NGFR}$ (Fig. 1) and undergo apoptotic cell death after serum withdrawal




Fig. 2. Enhancement of neural cell death in cells expressing $p75^{NGFR}$. CSM 14.1 cells (7) were grown in DMEM with FBS (10%) at 34°C and then switched to the restrictive temperature of 39°C for 36 hours. Cell death was then induced by replacement of the medium with serum-free DMEM [either alone or in combination with NGF (5 nM)], and cell viability was determined each day for 4 days. Viable cells were identified by trypan blue exclusion and by propidium iodide fluorescence. Differences between cells expressing $p75^{NGFR}$ and control cells were highly statistically significant ($P = 0.0001$ by two-way analysis of variance, $n = 5$, from four different stable transfections of each plasmid). Error bars represent standard deviations. (A) Effect of serum-free medium on viability of cells expressing $p75^{NGFR}$ in comparison to control cells. Squares, cell transfected with pBabe-puro- $p75^{NGFR}$; circles, cells transfected with pBabe-puro; triangles, cells transfected with pBabe-puro- $p75^{NGFR}$, grown in medium with 10% serum; diamonds, cells transfected with pBabe-puro, grown in medium with 10% serum. (B) Effect of NGF (5 nM) and monoclonal antibodies (10 μ g/ml) on cells expressing $p75^{NGFR}$ (closed bars) and control cells (open bars). Control mAb was directed against human p53 (anti-p53) (10 μ g/ml) (Pharmingen). Each pair showed a highly significant difference ($P < 0.01$ by paired t test, $n = 3$), except the mAb to $p75^{NGFR}$ (anti-p75) ($P < 0.05$) and the controls (no significant difference).

(12), were studied. In the presence of mAb binding to $p75^{NGFR}$ (10 μ g/ml), the number of cells undergoing cell death after serum withdrawal for 3 days was decreased from $78 \pm 8\%$ to $13 \pm 4\%$ ($P < 0.01$ by paired t test, $n = 3$), whereas the same concentration of control mAb did not affect cell survival. Furthermore, mutant PC12 cells lacking expression of $p75^{NGFR}$ (13) underwent very little cell death in serum-free medium ($12 \pm 6\%$ cell death after 3 days of serum-free medium, $n = 4$), whereas mutant PC12 cells derived in parallel (13) that retained expression of $p75^{NGFR}$ also retained the characteristic of undergoing cell death in response to serum withdrawal

($50 \pm 15\%$ cell death after 3 days of serum-free medium, $n = 4$; $P < 0.01$ by paired t test). As an additional control, another plasma membrane protein, β -amyloid precursor protein (β -APP₇₅₁), was expressed with the pBabe-puro expression vector in the same conditionally immortalized neural cell line (CSM 14.1), without effect on apoptosis (14). This does not exclude the possibility that the expression of other proteins may enhance apoptosis.

Although both NGF and mAb directed against $p75^{NGFR}$ enhanced cell survival, and although Trk A is not expressed by CSM 14.1 cells (Fig. 1A), it was possible that NGF inhibited the death of tempera-




Fig. 3. Apoptosis in R2 cells (10) transfected with pBabe-puro-p75^{NGFR}, but not in R2 cells transfected with pBabe-puro. Cells were grown in DMEM with FBS (10%) at 34°C, seeded at a density of 5×10^4 cells/cm², and placed at 39°C in serum-free medium. After 6 days in serum-free medium, propidium iodide was added at a concentration of 10 μ M, and cells were viewed with a Zeiss Axiovert microscope. (A and B) R2 cells transfected with pBabe-puro. (C and D) R2 cells transfected with pBabe-puro-p75^{NGFR}. (A and C) Phase contrast. (B and D) Fluorescence. In (D), many nuclei are fragmented, which is characteristic of apoptosis (single arrows mark some examples); other nuclei are homogeneously stained, also characteristic of apoptosis. The only example of a nonapoptotic nucleus in this field is denoted by a double arrow. Magnification, $\times 320$.

ure-sensitive immortalized neural cells expressing p75^{NGFR} by binding to the high-affinity receptor [dissociation constant (K_d) = 2.3×10^{-11} M (15)] rather than the low-affinity receptor [K_d = 1.7×10^{-9} M (15)]. Therefore, several concentrations of NGF were tested. The inhibition of cell death by NGF in this cell line was minimal at concentrations of NGF that bind only the high-affinity NGF receptor significantly (Fig. 4). In contrast, concentrations of NGF equaling or exceeding the affinity constant for binding to the low-affinity receptor increased cell survival (Fig. 4). Survival of control CSM 14.1 cells transfected with the expression construct lacking the p75^{NGFR} open reading frame was not increased by NGF (Fig. 4).

Thus, the expression of p75^{NGFR} resulted in an enhancement of neural cell death

in serum-free medium when p75^{NGFR} was not bound by ligand or antibody, whereas enhancement of survival beyond that of controls occurred with binding of the receptor. This dichotomous response defines a previously undescribed type of receptor function within the nervous system. This effect of p75^{NGFR} may account for the fact that some cells become dependent for their survival on the binding of NGF. Early neural cell precursors are independent of NGF, but during development specific neural cells become dependent on NGF (16). Increased expression of p75^{NGFR}, which has been shown to occur during development (17), could conceivably effect such a switch. Although binding of NGF to Trk A enhances cellular survival and differentiation (1), active induction of cell death in the absence of NGF may also occur, and




Fig. 4. Inhibition of conditionally immortalized neural cell death by various concentrations of NGF. CSM 14.1 cells were grown as described in Fig. 2. Serum-free medium included the indicated concentrations of NGF. Error bars represent standard deviations ($n = 3$).

this may be mediated at least in part by p75^{NGFR}. The type of cell death induced by p75^{NGFR}—apoptosis—is the same as that induced by growth factor withdrawal (18). However, we cannot exclude the possibility that p75^{NGFR} may under some conditions induce necrosis, especially because the TNFRs may mediate either apoptosis or necrosis (19). Our results suggest an additional function for p75^{NGFR} in neural cells, but have no bearing on the other functions ascribed to p75^{NGFR} or on the interaction of other neurotrophins, such as brain-derived neurotrophic factor, with p75^{NGFR}. However, the enhancement of neural cell survival by binding of NGF or mAb to p75^{NGFR} suggests that a similar effect might occur when p75^{NGFR} is bound by other neurotrophins. Neither do the results bear on the role of p75^{NGFR} in the death of non-neuronal cells, such as astrocytes or developing renal cells.

Somewhat similar receptors have been described, including the TNFRs, FAS (Apo-1), and CD40. These molecules show general structural similarity to p75^{NGFR}, with similar extracellular cysteine-rich pseudo-repeats and a single transmembrane domain (20). The structural similarity of p75^{NGFR} to the other members of the superfamily occurs in the extracellular domain (5), but the functional similarity may result from the transduction of a signal leading to (or inhibiting) cell death. The function of p75^{NGFR} is analogous to that of CD40 in that expression occurs on developing cells [mainly central cholinergic, sympathetic, and sensory neurons in the case of p75^{NGFR}, centroblasts and centrocytes in the case of CD40 (6)], and leads to a requirement for binding if survival is to occur. In both cases, binding of the receptor leads to improved, but incomplete, cell survival (Figs. 2 and 4) (6). Other determinants are clearly involved, because binding of antigen by developing B cells also enhances survival (6), lack of expression of CD40 ligand does not result in a reduction

in circulating B cells (21), and neural cells expressing p75^{N^{GR}} survive in media containing serum (Fig. 2). The mechanism by which unbound p75^{N^{GR}} or other members of this receptor superfamily lead to neural cell death is unknown. However, the structural and functional relation between p75^{N^{GR}} and TNFR I and II suggests that they may have similar mechanisms of action.

The highest level of expression of p75^{N^{GR}} in the central nervous system occurs in cholinergic neurons of the nucleus basalis of Meynert, the cells most severely affected in Alzheimer's disease. These cells continue to express normal (22) or supra-normal (23) amounts of p75^{N^{GR}} mRNA and protein during the neuronal degeneration associated with Alzheimer's disease. In contrast, cholinergic cells of the brainstem that resemble those of the nucleus basalis morphologically, but do not express p75^{N^{GR}} (24), do not degenerate in Alzheimer's disease (25).

REFERENCES AND NOTES

1. R. Klein, S. Jing, V. Nandori, E. O'Rourke, M. Barbacid, *Cell* 66, 189 (1991); B. L. Hempstead, D. Martin-Zanca, D. R. Kaplan, L. F. Parada, M. V. Chao, *Nature* 350, 678 (1991); C. F. Ibanez et al., *Cell* 69, 329 (1992).
2. N. Y. Ip et al., *Neuron* 10, 137 (1993).
3. T. J. Schall et al., *Cell* 61, 361 (1990).
4. C. A. Smith et al., *Science* 248, 1019 (1990).
5. B. C. Trauth et al., *ibid.* 245, 301 (1989); N. Itoh et al., *Cell* 66, 233 (1991).
6. Y. J. Liu et al., *Nature* 342, 21 (1989); J. Blancher, P. de Paoli, A. Valté, E. Garcia, F. Roussel, *Science* 251, 70 (1991).
7. M. Durand, D. C. Chugani, M. Mahmoudi, M. E. Phelps, *Soc. Neurosci. Abstr.* 16, 40 (1990).
8. J. P. Morgenstern and H. Land, *Nucleic Acids Res.* 18, 3587 (1990).
9. L. T. Zhong, S. P. Mah, R. H. Edwards, D. E. Brøderick, *Soc. Neurosci. Abstr.* 18, 44 (1992); L. T. Zhong et al., *Proc. Natl. Acad. Sci. U.S.A.* 90, 4539 (1993).
10. M. E. Greenberg, R. Brackenbury, G. M. Edelman, *Proc. Natl. Acad. Sci. U.S.A.* 81, 969 (1984).
11. J. F. R. Kerr and B. V. Harmon, in *Apoptosis: The Molecular Basis of Cell Death*, L. D. Tomei and F. O. Cope, Eds. (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1991), vol. 3, p. 321.
12. A. Rukenstein, R. E. Rydel, L. A. Greene, J. Neurosci. 11, 2552 (1991); S. P. Mah et al., *J. Neurochem.* 60, 1183 (1993).
13. PC12 cells were maintained in Dulbecco's minimum essential medium (DMEM; Gibco) containing 5% horse serum and 5% supplemented calf serum (HyClone), in 12% CO₂ at 37°C. Cells were mutagenized with 10 mM ethyl methanesulfonate for 6 hours and then washed with 2 x 10 ml of medium. After growing for 10 days, cells were trypsinized and subcultured into 36 plates at an approximate density of 10⁴ cells per plate. The cells were grown for 10 days (approximately five divisions) and then treated with NGF (25 ng/ml) for 5 days. Groups of clonally derived cells that did not respond to NGF by extending neurites were isolated, subcloned, and then tested for the absence of NGF-induced neurite outgrowth and the presence (or absence) of p75^{N^{GR}} by Northern (RNA) and protein immunoblot analysis. The two subclones used in these experiments were the NFA5 subclone, which does not express p75^{N^{GR}}, and NRS0, which does express p75^{N^{GR}}.
14. S. Rostkovich et al., unpublished data.
15. A. Suter, R. J. Riopelle, R. M. Harris-Warrick, E.

- Shooter, *J. Biol. Chem.* 254, 5972 (1979).
16. R. Levi-Montalcini, *Harvey Lect.* 60, 217 (1986); *Science* 237, 1154 (1987).
17. T. H. Large et al., *Neuron* 2, 1129 (1989).
18. D. P. Martin et al., *J. Cell Biol.* 106, 829 (1988).
19. K. Schulte-Osthoj et al., *J. Biol. Chem.* 267, 5317 (1992).
20. N. Itoh et al., *Cell* 66, 233 (1991).
21. J. P. DiSanto, J. Y. Bonnet, J. F. Gauchat, A. Fischer, G. de Saint Basille, *Nature* 361, 841 (1993).
22. M. Goedert, A. Fine, D. Dawbarn, G. K. Wilcock, M. V. Chao, *Mol. Brain Res.* 5, 1 (1989); J. H. Kordower, D. M. Gash, M. Bothwell, L. Herth, E. J. Mufson, *Neurobiol. Aging* 10, 67 (1989).
23. P. Emfors, N. Lindfors, V. Chan-Palay, H. Persson, *Dementia* 1, 138 (1990).
24. N. J. Woolf, E. Gould, L. L. Butcher, *Neuroscience* 30, 143 (1989).
25. N. J. Woolf, R. W. Jacobs, L. L. Butcher, *Neurosci. Lett.* 68, 277 (1989).
26. P. Chomczynski and N. Sacchi, *Anal. Biochem.* 162, 156 (1987).
27. S. P. Mah et al., *J. Neurochem.* 60, 1183 (1993).
28. The p75^{N^{GR}} cDNA in pUC9 was digested with Sal I, filled in with Klenow fragment and deoxynucleotide triphosphates, and then digested with Bgl II. The 1.7-kb fragment containing the entire open reading frame of p75^{N^{GR}} was then ligated into pUC18 that had been digested with Sma I and Bam HI. The resulting plasmid was digested with Eco 47III and Sal I and ligated into pBabe-puro (8) that had been cut with Sna BI and Sal I, to create pBabe-puro-p75^{N^{GR}}. CSM 14.1 cells (7) are rat nigral neural cells immortalized with the temperature-sensitive large T antigen of SV40. These cells express tyrosine hydroxylase, neuron-specific enolase, and neurofilament (NF-L). CSM 14.1 cells were transfected with pBabe-puro-p75^{N^{GR}} with the cationic lipid DOTAP (Boehringer-Mannheim, Inc.) and then selected in puromycin (7 µg/ml). The comparison of single colonies

can introduce bias into the results (9), but this was obviated by comparison of entire pools of stable transfectants (9); therefore, pools of stable transfectants (populations including more than 100 separate colonies) with pBabe-puro-p75^{N^{GR}} were compared with pools of pBabe-puro transfectants. Cells were grown in DMEM with fetal bovine serum (FBS) (10%) at 34°C in 5% CO₂. Total RNA was prepared by the method of Chomczynski (26), and electrophoresis was carried out in formaldehyde gels. After Northern transfer to nylon, ³²P-labeled probes for p75^{N^{GR}} (1-kb cDNA fragment, digested with Stu I), Trk A (0.5-kb cDNA fragment, digested with Xho I), and γ-actin were hybridized sequentially. Blots were exposed to film for 24 hours for the p75^{N^{GR}} and Trk A probes and for 2 hours for the γ-actin probe. For immunocytochemistry, cells were fixed in paraformaldehyde (4%) for 15 min and permeabilized in 0.1% Triton X-100. Immunocytochemistry was done as described (27), with a polyclonal antibody (1:2500) to purified p75^{N^{GR}}. As controls, primary antibody was omitted and control transfectants were stained; both of these controls showed a similar lack of staining.

29. M. J. Radake, T. P. Misko, C. Hsu, L. A. Herzenberg, E. M. Shooter, *Nature* 325, 593 (1987).
30. We thank E. Shooter for the p75^{N^{GR}} cDNA and hybridoma cells secreting the 192 immunoglobulin G mAb, G. Westkamp for the polyclonal antibody to p75^{N^{GR}}, H. Land for the pBabe-puro expression vector, M. Cohen for the F2 cells, M. Durand for the CSM14.1 cells, E. Koo for the APP^Δ cDNA, T. Ord for technical assistance, and R. Edwards and B. Howard for thoughtful discussions. Supported by a pilot grant from the Alzheimer's Disease and Related Disorders Association, NIH grant AG10671 (to D.E.B.), and NIH grant NS10328 (to L.L.B.).

16 February 1993; accepted 3 May 1993

Redundant Mechanisms of Calcium-Induced Calcium Release Underlying Calcium Waves During Fertilization of Sea Urchin Eggs

Antony Gallone,* Alex McDougall, William B. Busa, Nick Willmott, Isabelle Gillot, Michael Whitaker

Propagating Ca²⁺ waves are a characteristic feature of Ca²⁺-linked signal transduction pathways. Intracellular Ca²⁺ waves are formed by regenerative stimulation of Ca²⁺ release from intracellular stores by Ca²⁺ itself. Mechanisms that rely on either inositol triphosphate or ryanodine receptor channels have been proposed to account for Ca²⁺ waves in various cell types. Both channel types contributed to the Ca²⁺ wave during fertilization of sea urchin eggs. Alternative mechanisms of Ca²⁺ release imply redundancy but may also allow for modulation and diversity in the generation of Ca²⁺ waves.

Transient increases in the concentration of calcium ions ([Ca²⁺]_i) act as cell signals. In general, the signal shows spatial and temporal inhomogeneity and takes the form of waves or oscillations within the cell (1).

A. Gallone and N. Willmott, Department of Pharmacology, Oxford University, Oxford OX1 3QT, United Kingdom.

A. McDougall, I. Gillot, M. Whitaker, Department of Physiology, University College London, London WC1E 6BT, United Kingdom.

W. B. Busa, Department of Biology, Johns Hopkins University, Baltimore, MD 21218.

*To whom correspondence should be addressed.

Several mechanisms have been proposed to account for regenerative Ca²⁺ release (2). Release of Ca²⁺ from internal stores can be stimulated by an increase in [Ca²⁺]_i; this process is termed Ca²⁺-induced Ca²⁺ release (CICR) (3). This Ca²⁺ release appears to be mediated by Ca²⁺ channels in the endoplasmic reticulum (ER) that are sensitive to cytoplasmic agonists, to [Ca²⁺]_i, and to the amount of Ca²⁺ in the lumen of the ER (4). Two closely related Ca²⁺ channels with these properties are the inositol triphosphate (IP₃) receptor (IP₃R) (5) and the